

PLASMA BUBBLES AND EQUATORIAL SPREAD F IRREGULARITY STUDIES OVER BRAZIL

I. S. Batista, M. A. Abdu, E. R. de Paula, J. H. A. Sobral,
P. Muralikrishna and R. T. de Medeiros¹

Instituto Nacional de Pesquisas Espaciais
Cx. Postal 515, São José dos Campos, SP, Brazil 12201-970

ABSTRACT

For more than 20 years ionospheric studies have been conducted in Brazil, in order to understand the plasma bubble and spread F irregularity phenomena and their influence on radio-wave propagation. Most of the studies were carried out using ground-based equipments (ionosondes, polarimeters, scintillation measurements and photometers), operating in Cachoeira Paulista (22.5°S, 45°W), São José dos Campos (23°S, 46°W) and Fortaleza (4°S, 38°W). In situ measurements carried out using payloads on board of sounding rockets launched from Natal (5.9°S, 35.3°W) have also contributed to the study of the plasma bubble phenomena over Brazil. The equatorial spread F phenomena are often associated with plasma depleted magnetic flux tubes, namely, plasma bubbles. The plasma bubbles are generated over the equatorial F region through the Rayleigh-Taylor gravitational instability process. The enhanced post-sunset $E \times B$ drift is of prime importance on the generation of plasma bubbles and equatorial spread F irregularities. The magnetic declination angle over Brazil is very high (it attains a global maximum over Brazilian longitude sector). The observational studies and theoretical simulations conducted in Brazil have shown that the very high magnetic declination angle greatly affects the seasonal distribution of spread F occurrence. In this paper, all these observational aspects as well as the simulations conducted to explain them, are reviewed.

INTRODUCTION

The plasma bubble phenomena are regions of depleted plasma that occurs in the low and equatorial latitude nighttime ionospheric F-region. They are generated by plasma instability processes that set in under the ambient ionospheric conditions governed by sunset electrodynamics, the most obvious manifestation of which is the rapid vertical uplift of the F-layer. The primary instability is believed to be driven by the Rayleigh-Taylor (R-T) gravitational fluid instability process. The steeping density gradients that results from the primary instability mechanism then become unstable to secondary instability processes leading to the generation of smaller scale sizes in a hierarchy of irregularities. Basically, the thermospheric zonal wind that blows eastward across the sunset terminator is responsible for the F layer dynamo electric field. In the presence of a longitudinal gradient in the F-layer Pedersen conductivity, this results in the generation of eastward electric field which contributes to the vertical uplift of the layer and to the instability growth rate factor (γ).

GROUND BASED EQUIPMENTS

Plasma bubble manifest themselves in ionograms as diffuse echoes known as spread F. The ionosondes operating in Fortaleza (FZ) and Cachoeira Paulista (CP) have given a great contribution to the understanding of the plasma bubble morphology and spread F characteristics over the Brazilian region [1-13]. The analysis of simultaneous data from FZ and CP permitted the study of the plasma bubble vertical rise velocities [7], their seasonal and solar cycle dependences [5, 10] and the important aspect of the magnetic declination control of equatorial spread F and F region dynamo [2, 8, 9, 11, 13]. The spread F associated irregularities, once developed after sunset (1830-1930 LT) go through a statistical maximum during the premidnight hours (~22 LT) with a decay phase that could continue till sunrise (~6

¹On leave from Universidade Federal do Rio Grande do Norte

LT). The onset local times at locations away from the magnetic equator get later with increasing latitude due to the fact that the depleted flux tubes rising upward over the equator are responsible for extending the ESF belt to latitudes away from the equator in such a way that the width of the belt is defined by the flux tube extremities (feet) intersections with the low latitude F regions. This is also a reason for the statistical duration of the spread F over low latitude to be somewhat shorter than over the equator. Similar statistics should apply for VHF/UHF scintillations as well [14-17]. The use of spaced VHF receptor have allowed to infer the plasma bubble zonal velocities over low latitudes in Brazil [16,17].

Since 1981 scanning photometer data have been used to study ionospheric plasma bubbles. Their manifestation in the airglow measurements are depletions in 630 nm airglow profiles over CP [18-26]. They have been used as tracers of the zonal plasma drift velocities [18,24] and the results are comparable to the obtained by VHF polarimeters [24].

IN SITU MEASUREMENTS

In situ measurements of electron density height profile of the equatorial nighttime ionosphere, under a developing spread F event was carried out using a high frequency capacitance probe on board of a sounding rocket launch from Natal, Brazil [27]. During the upleg of the flight the experiment detected a series of plasma bubbles in varying degree of their growth phase. The launch criteria, based on ionosonde and scintillation data (no radar was available to the campaign), supported the view that the range type spread F, observed in the ionogram over low latitude, are indeed produced by flux tube aligned depletion that developed upward over the equator under the gravitational R-T instability mechanism. Another rocket experiment was conducted from Natal, this time to inject chemical components in the bottomside equatorial F2 region, to create a hole in ionization [28]. The hole induced depletions were observed in total electron content (TEC) monitoring stations and in ionosondes located 300 km magnetically east of the chemical release point.

The satellite experiments give a worldwide coverage of plasma bubble and spread F irregularities occurrence. The results from the DE2 and AE satellite [29-31] gave information about zonal and vertical plasma drifts (electric fields) in the low latitude F region and its seasonal variation. The BSS (bottomside sinusoidal irregularities) have been identified in satellite measurements [32-34] and is believed to be correlated with the frequency type spread F seen in low latitude ionograms. The Japanese satellites have given a great contribution to the global studies of spread F and plasma bubble occurrence and characteristics [35-38].

MODELS

In order to explain the seasonal and longitudinal variation on spread F occurrence over Brazilian region, when compared with Jicamarca, Peru, a numerical model was developed [13]. That model took in account the effect of the magnetic declination over the F region dynamo development, that was essential to explain most of the peculiar features observed over the Brazilian region [13, 39, 40]. Although this model self-consistently solves the E and F region electrodynamics, their electron density profile (input) is a very simple Chapman layer and needs improvement.

CONCLUSIONS

The studies conducted so far have brought out a fairly good description of the morphology of the irregularities. Good progress has been achieved in our understanding of the seasonal/longitudinal effects. There are strong indications that the day-to-day changes in the thermospheric wind (both magnitude and direction) could be the major key to explain the

ESF variability. Therefore, systematic and regular measurements should be carried out at key locations in the equatorial regions.

REFERENCES

1. Batista, I. S.; de Paula, E. R.; Abdu, M. A.; Trivedi, N. B.; Greenspan, M. E. Ionospheric effects of the March 13, 1989, magnetic storm at low and equatorial latitudes. *J. Geophys. Res.*, **96**, 13943-13952, 1991.
2. Abdu, M. A.; Batista, I. S.; Sobral, J. H. A. A new aspect of magnetic declination control of equatorial spread F and F region dynamo. *J. Geophys. Res.*, **97**, 14897-14904, 1992.
3. Medeiros, R. T.; Abdu, M. A.; Kantor, I. J. A comparative study of VHF scintillation and spread F events over Natal and Fortaleza in Brazil. *J. Geophys. Res.*, **88**, 6253-6258, 1983.
4. Abdu, M. A.; Medeiros, R. T.; Sobral, J. H. A. Equatorial spread F instability conditions as determined from ionograms. *Geophys. Res. Lett.*, **9**, 692-695, 1982.
5. Abdu, M. A.; Medeiros, R. T.; Bittencourt, J. A.; Batista, I. S. Vertical ionization drift velocities and range type spread F in the evening equatorial ionosphere. *J. Geophys. Res.*, **88**, 399-402, 1983.
6. Abdu, M. A.; Medeiros, R. T.; Nakamura, Y. Latitudinal and magnetic flux tube extension of the equatorial spread F irregularities. *Geophys. Res. Lett.*, **88**, 4861-4868, 1983.
7. Abdu, M. A.; Medeiros, R. T.; Sobral, J. H. A.; Bittencourt, J. A. Spread F plasma bubble vertical rise velocities determined from spaced ionosonde observations. *J. Geophys. Res.*, **88**, 9197-9204, 1983.
8. Abdu, M. A.; Bittencourt, J. A.; Batista, I. S. Magnetic declination control of the equatorial F-region dynamo electric field development and spread-F. *J. Geophys. Res.*, **86**, 11443, 1981.
9. Abdu, M. A.; Batista, I. S.; Bittencourt, J. A. Some characteristics of spread F at magnetic equatorial station Fortaleza. *J. Geophys. Res.*, **86**, 6836-6842, 1981.
10. Abdu, M. A.; Sobral, J. H. A.; Nelson, O. R.; Batista, I. S. Solar cycle related range type spread-F occurrence characteristics over equatorial and low latitude stations in Brazil. *J. Atmos. Terr. Phys.*, **47**, 901-905, 1985.
11. Abdu, M. A.; Bittencourt, J. A.; Batista, I. S. Longitudinal differences in the spread F characteristics. *Rev. Bras. Geof.*, **13**, 647-664, 1983.
12. Nelson, O. R.; Abdu, M. A.; Batista, I. S. Equatorial F-region irregularity latitudinal extension and dynamo electric field. *J. Atmos. Terr. Phys.*, **48**, 181-186, 1986.
13. Batista, I. S.; Abdu, M. A.; Bittencourt, J. A. Equatorial F region vertical plasma drifts: Seasonal and longitudinal asymmetries in the American sector. *J. Geophys. Res.*, **91**, 12055-12064, 1986.
14. Yeh, K. C.; Soicher, H.; Liu, C. H.; Bonelli, E. Ionospheric bubbles observed by the Faraday rotation method at Natal, Brazil. *Geophys. Res. Lett.*, **6**, 473-475, 1979.
15. Yeh, K. C.; Mullen, J. P.; Medeiros, J. R.; Silva, R. F.; Medeiros, R. T. Ionospheric scintillation observed at Natal. *J. Geophys. Res.*, **86**, 7527-7532, 1981.
16. Abdu, M. A.; Kantor, I. J.; Batista, I. S.; de Paula, E. R. East-west plasma bubble irregularity motion determined from spaced VHF polarimeters: Implication on velocity shear in the zonal F region bulk plasma motion. *Radio Sci.*, **20**, 111-122, 1985.
17. Abdu, M. A.; Batista, I. S.; Sobral, J. H. A.; de Paula, E. R.; Kantor, I. J. Equatorial ionospheric plasma bubble irregularity occurrence and zonal velocities under quiet and disturbed conditions, from polarimeter observations. *J. Geophys. Res.*, **90**, 9921-9928, 1985.
18. Sobral, J. H. A.; Abdu, M. A. Latitudinal gradient in the plasma bubble zonal velocities as observed by scanning 630-nm airglow measurements. *J. Geophys. Res.*, **95**, 8253-8257, 1990.
19. Abdu, M. A.; Sobral, J. H. A.; Nakamura, Y. On the low latitude scanning photometer signatures of the equatorial ionosphere plasma bubbles. *Rev. Bras. Geofis.*, **3**, 33-39, 1985.
20. Sobral, J. H. A.; Abdu, M. A.; Zamlutti, C. J.; Batista, I. S. Association between plasma bubble irregularities and airglow disturbances over Brazilian low latitudes. *Geophys. Res. Lett.*, **7**, 980-982, 1980.
21. Sobral, J. H. A.; Abdu, M. A.; Batista, I. S. Airglow studies on the ionosphere dynamics over low latitude in Brazil. *Ann. Geophys.*, **36**, 199-204.
22. Sobral, J. H. A.; Abdu, M. A. Solar activity effects on equatorial plasma bubble zonal velocity and its latitude gradient as measured by airglow scanning photometers. *J. Atmos. Terr. Phys.*, **8**, 729-742, 1991.

23. Sobral, J. H. A.; Abdu, M. A.; Batista, I. S.; Zamlutti, C. J. Wave disturbances in the low latitude ionosphere and equatorial ionospheric plasma depletions. *J. Geophys. Res.*, 86, 1374-1378, 1981.
24. Abdu, M. A.; Sobral, J. H. A.; Nakamura, Y.; Zamlutti, C. J. Equatorial plasma bubble zonal velocity height gradient from spaced VHF polarimeter and scanning 630NM measurements. *Geophys. Res. Lett.*, 14, 965-968, 1987.
25. Bittencourt, J. A.; Sahai, Y. Low-latitude ionospheric dynamical processes from atomic oxygen nightglow emissions. *Rev. Bras. Geofis.*, 9, 85-93, 1991.
26. Nakamura, Y.; Sobral, J. H. A.; Abdu, M. A. A simulation of equatorial plasma bubble signatures on the OI 6300 Å nightglow meridional profile over Brazilian low latitude. *Rev. Bras. Geofis.*, 2, 65-77, 1984.
27. Klobuchar, J. A.; Abdu, M. A. Equatorial ionosphere irregularities Produced by the Brazilian ionospheric modification experiment (BIME). *J. Geophys. Res.*, 94, 2721-2726, 1989.
28. Abdu, M. A.; Muralikrishna, P.; Batista, I. S.; Sobral, J. H. A. Rocket observation of the equatorial plasma bubbles over Natal, Brazil. using a High-Frequency Capacitance Probe. *J. Geophys. Res.*, 96, 7689-7695, 1991.
29. Maynard, N. C.; Aggson, T. L.; Herrero, F. A.; Liebrecht, M. C. Average low-latitude meridional electric fields from DE 2 during solar maximum. *J. Geophys. Res.*, 93, 4021-4037, 1988.
30. Coley, W. R.; Heelis, R. A. low-latitude zonal and vertical ion drifts seen by DE 2. *J. Geophys. Res.*, 94, 6751-6761, 1989.
31. Dachev, Ts. P.; Walker, J. C. G. Seasonal dependence of the distribution of large-scale plasma depletions in the low-latitude F region. *J. Geophys. Res.*, 87, 7625-7639, 1982.
32. Cragin, B. L.; Valladares, C. E.; Hanson, W. B.; McClure, J. P. Bottomside sinusoidal irregularities in the equatorial F region 2. Cross-correlation and spectral analysis. *J. Geophys. Res.*, 90, 1721-1734, 1985.
33. Valladares, C. E.; Hanson, W. B.; McClure, J. P.; Cragin, B. L. Bottomside sinusoidal irregularities in the equatorial F region. *J. Geophys. Res.*, 88, 8025-8042, 1983.
34. Basu, S.; Basu, S. Scintillation associated with bottomside sinusoidal irregularities in the equatorial F region. *J. Geophys. Res.*, 91, 270-276, 1986.
35. Maruyama, T.; Matuura, N. Longitudinal variability of annual changes in activity of equatorial spread F and plasma bubbles. *J. Geophys. Res.*, 89, 10903-10912, 1984.
36. Oya, H.; Takahashi, T.; Watanabe, S. Observation of low latitude ionosphere by the impedance probe on board the Hinotori satellite. *J. Geomag. Geoelectr.*, 38, 111-123, 1986.
37. Watanabe, S.; Oya, H. Occurrence characteristics of low latitude ionospheric irregularities observed by impedance probe on board the Hinotori satellite. *J. Geomag. Geoelectr.*, 38, 125-149, 1986.
38. Takahashi, T.; Oya, H.; Watanabe, S. Ionospheric disturbances induced by substorm associated electric fields in the low-latitude F-region. *J. Geomag. Geoelectr.*, 39, 187-209, 1987.
39. Batista, I. S.; Abdu, M. A.; Medrano, R. A. Magnetic Activity effects on range type spread-F and vertical plasma drifts at Fortaleza and Huancayo as studied through ionosonde measurements and theoretical modeling. *Ann. Geophys.*, 8, 357-364.
40. Batista, I. S.; Abdu, M. A. Simulação numérica das componentes zonais da deriva do plasma e do vento termosférico em regiões equatoriais. *Rev. Bras. Geofis.*, 4, 135-141, 1986.