

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/319007216>

Gross Primary Productivity in the northern region of Para state, Brazilian amazon, from MOD17 data

Conference Paper · July 2017

CITATIONS

0

READS

38

7 authors, including:



Nathaniel A Brunsell

University of Kansas

115 PUBLICATIONS 1,425 CITATIONS

SEE PROFILE



Gabriel Bertani

National Institute for Space Research, Brazil

13 PUBLICATIONS 2 CITATIONS

SEE PROFILE



Thiago Veloso dos Santos

University of Minnesota Twin Cities

7 PUBLICATIONS 19 CITATIONS

SEE PROFILE



Luiz E. O. C. Aragão

National Institute for Space Research, Brazil

225 PUBLICATIONS 6,610 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



PAST: Pre-Columbian Amazon Scale Transformations [View project](#)



SOYBEAN YIELD ESTIMATES BASED ON TEMPORALLY STABLE PIXELS USING MODIS/EVI DATA [View project](#)

All content following this page was uploaded by [Thiago Veloso dos Santos](#) on 09 August 2017.

The user has requested enhancement of the downloaded file.

GROSS PRIMARY PRODUCTIVITY IN THE NORTHERN REGION OF PARA STATE, BRAZILIAN AMAZON, FROM MOD17 DATA

Gabriel de Oliveira^{1,2}, *Nathaniel A. Brunsell*¹, *Elisabete C. Moraes*², *Yosio E. Shimabukuro*², *Gabriel Bertani*², *Thiago V. dos Santos*³, *Luiz E. O. C. Aragao*²

Department of Geography and Atmospheric Science, University of Kansas, Lawrence, KS 66045, US; Remote Sensing Division, National Institute for Space Research, Sao Jose dos Campos, SP 12227-010, Brazil; Department of Soil, Water and Climate, University of Minnesota, Saint Paul, MN 55108, US
E-mail: gabrieloliveira@ku.edu

ABSTRACT

This study aimed to characterize and analyze, based on MOD17 data, the spatio-temporal dynamics of GPP in the northern region of Para state, Brazilian Amazon, during a 6-year period (2001 to 2006). The study area encompasses two river basins (Upper Tapajos and Curua-Una) and covers ~74,190 km². The spatial variation of GPP was primarily related to the larger presence of forested areas in Upper Tapajos River basin in comparison with Curua-Una River basin. Temporally, GPP varied with the dry and wet seasons in the region. There was a decrease of ~4% in GPP during the dry season, which was related to the fact that the water limitation during the dry season in Amazonia leads to a decrease in photosynthesis, affecting vegetation productivity. It was observed a reasonable interannual variation of GPP in the study area, which corresponded to ~10%.

Index Terms - vegetation productivity, photosynthesis, Amazonia, tropical forest, MOD17.

1. INTRODUCTION

The Amazonian forest has a remarkable influence on climate primarily through by the release or uptake of atmospheric CO₂ [1], [2]. As the initial carbon fixed by vegetation through photosynthesis, gross primary productivity (GPP) is a key driver of the global carbon cycle [3]. In this context, knowing the annual and interannual variability of gross primary productivity (GPP) in the Amazon ecosystem is critical for predicting how it will respond to future physical and biological perturbations.

The purpose of this study was to characterize and analyze, based on MOD17 product, the spatio-temporal dynamics of GPP in the northern region of Para state, Brazilian Amazon, during a 6-year period (January 2001 to December 2006).

2. MATERIALS AND METHODS

2.1. Study area

The study area is located in the northern region of the state of Para, Brazil. The study area encompasses two river basins (Upper Tapajos and Curua-Una) and covers ~74,190 km² (Fig. 1). The climate according Köppen classification is the AmW, with an annual mean precipitation of ~2000 mm and an annual mean air temperature of ~26 °C. The wet season is from January to June and the dry season lasts from July through December [4]. The main natural vegetation type in the study area is the dense forest [5]. Due to deforestation, it is possible to observe extensive patches of different land cover types such as pasture, secondary succession forest, etc.

2.2. MOD17 data

The MOD17 product [6] is calculated based on the radiation-use efficiency (RUE) concept [7]:

$$GPP = \varepsilon f(T_{ar,min}) f(VPD) APAR \quad (1)$$

where ε is the photosynthetically active radiation (PAR) conversion efficiency; $f(T_{ar,min})$ and $f(VPD)$ are scale factors related to minimum air temperature and vapour pressure deficit, respectively. The algorithm is driven by meteorological information derived from reanalysis (e.g. solar radiation and vapour pressure deficit), and biophysical vegetation parameters retrieved through remote sensing (e.g. fraction of photosynthetically active radiation (fPAR) and leaf area index (LAI)).

We used the product MOD17A2, which provides GPP in a monthly composite at 1 km resolution. The images used encompassed the h12v09 tile, which covers the Upper Tapajos and Curua-Una River basins in the Amazon region, from January to December, over 6 years (2001-2006). First, MODIS images were projected to a geographic coordinate system (lat./long.) based on the WGS 84 datum and converted to the GeoTIFF format using the MODIS Reprojection Tool (MRT). Then, a number of steps were

undertaken using Python Language Reference 2.7. These steps included clipping of the study area, multiplication by

scale factors, and application of the land-water mask extracted from product MOD44W over the dataset.




Fig. 1. Map showing the location of the study area.




Fig. 2. Monthly averages of GPP ($\text{g C m}^{-2} \text{ month}^{-1}$), between the years of 2001 and 2006.

3. RESULTS AND DISCUSSIONS

Fig. 2 presents the spatial distribution of the monthly average GPP (January to December), from 2001 and 2006, in the Upper Tapajós and Curua-Una River basins. The images from January to August presented higher values of GPP in comparison with the images from September to December. This pattern is related to the characteristics of the dry and wet seasons in the eastern part of Amazonia. The dry season can vary interannually, but typically occurs between July and December. The wet season typically occurs between January and June. The degree of water limitation predicts seasonal patterns of photosynthesis in the southern Amazon. The low water availability and increased vapour pressure deficit during the dry period induces stomatal closure, which leads to a decline in the photosynthesis rate [8]. Thus, the decrease of GPP in the dry season occurred as expected and it is corroborated by several studies conducted in the region [2], [9]. The highest values of GPP are shown to be situated in the western part of the study area (Upper Tapajós river basin), which has experienced less deforestation compared with the eastern

region (Curua-Una river basin). It is interesting to note in the cyan dashed circles over the images of June, July, August and September the spatial pattern of deforestation in the Amazon, known as the fish bone. It is possible to verify the difference of GPP values between logged and forested areas, with the later presenting higher values. This is because herbaceous ecosystems are less productive than forest ecosystems [10].

The monthly GPP ranged between 103.2 and 263.0 g C m⁻² month⁻¹, representing an absolute difference of 159.8 g C m⁻² (Fig. 3). These values were observed in September 2006 and July 2005, respectively. On average for the whole period, the monthly GPP was 193.4 g C m⁻² month⁻¹. The monthly average GPP varied between 158.1 and 246.0 g C m⁻² month⁻¹. November presented the lowest values of GPP while July presented the highest ones. The monthly average GPP in the wet season was 197.5 g C m⁻² month⁻¹ while in the dry season it was 189.3 g C m⁻² month⁻¹, which means a decrease in GPP during the dry season of ~4%. The annual GPP values ranged from 2223.7 (2005) to 2446.3 g C m⁻² year⁻¹ (2003), which corresponds to a variation of ~10%.




Fig. 3. Monthly GPP between January 2001 and December 2006 over the study area. The dashed orange line represents the moving average of time series (period=2).

4. CONCLUDING REMARKS

The use of MOD17 data enabled mapping the spatial distribution and evaluating GPP in the northern region of Para state, as well as analyzing the temporal dynamics of this variable in Amazon rainforest remnants for the period between 2001 and 2006. The spatial variation of GPP was primarily related to the larger presence of forested areas in Upper Tapajós River basin (western part) in comparison with Curua-Una River basin (eastern part), which is situated

within the so-called arc of deforestation. Temporally, GPP varied with the dry and wet seasons in the region. There was a decrease in GPP during the dry season, corresponding to ~4%. It occurred as expected and it was related, among others, to the fact that the water limitation during the dry season in Amazonia leads to a decrease in photosynthesis, affecting vegetation productivity. It was observed a reasonable interannual variation of GPP in the study area, which corresponded to ~10%.

5. ACKNOWLEDGEMENT

Gabriel de Oliveira acknowledges CNPq (Grant No. 52521/2012-7) and CAPES (Grant No. 8210/2014-4) agencies for providing research fellowships. Luiz E. O. C. Aragao acknowledges the financial support of FAPESP

(Grant No. 50533-5) and CNPq (Grant No. 304425/2013-3) agencies.

6. REFERENCES

[1] M.H. Costa, and J.A. Foley, "Combined effects of deforestation and doubled atmospheric CO₂ concentrations

on the climate of Amazonia”, *J. Clim.*, vol. 13, n.1, pp. 18-34, 2000.

[2] J.E. Lee, C. Frankenberg, C. van der Tol, J.A. Berry, L. Guanter, C.K. Boyce, J.B. Fisher, E. Morrow, J.R. Worden, S. Asefi, and G. Badgley, “Forest productivity and water stress in Amazonia: Observations from GOSAT chlorophyll fluorescence”, *Proc. R. Soc. Lond. B. Biol. Sci.*, vol. 280, no. 1761, pp. 1-9, 2013.

[3] Y. Zhang, C. Song, G. Sun, L.E. Band, S. McNulty, A. Noormets, Q. Zhang, and Zhang, Z., “Development of a coupled carbon and water model for estimating global gross primary productivity and evapotranspiration based on eddy flux and remote sensing data”. *Agric. For. Meteorol.*, vol. 223, pp. 116-131, 2016.

[4] S.R. Saleska, S.D. Miller, D.M. Matross, M.L. Goulden, S.C. Wofsy, H.R. Da Rocha, P.B. De Camargo, P. Crill, B.C. Daube, H.C. De Freitas, and L. Hutyyra, “Carbon in Amazon forests: unexpected seasonal fluxes and disturbance-induced losses”, *Science*, vol. 302, no. 5650, pp. 1554-1557, 2003.

[6] M. Zhao, F.A. Heinsch, R.R. Nemani, and S.W. Running, “Improvements of the MODIS terrestrial gross and net primary production global data set”, *Remote Sens. Environ.*, vol. 95, pp. 164-176, 2005.

[7] J. L. Monteith, "Solar radiation and productivity in tropical ecosystems." *J. Appl. Ecol.*, vol. 9, no. (3), pp. 747-766, 1972.

[8] N. Restrepo-Coupe, H.R. da Rocha, L.R. Hutyyra, A.C. da Araujo, L.S. Borma, B. Christoffersen, O.M. Cabral, P.B. de Camargo, F.L. Cardoso, A.C.L. da Costa, and D.R. Fitzjarrald, “What drives the seasonality of photosynthesis across the Amazon basin? A cross-site analysis of eddy flux tower measurements from the Brasil flux network”, *Agric. For. Meteorol.*, vol. 182, pp. 128-144, 2013.

[9] L.R. Hutyyra, J.W. Munger, S.R. Saleska, E. Gottlieb, B.C. Daube, A.L. Dunn, , D.F. Amaral, , P.B. De Camargo, and S.C. Wofsy, “Seasonal controls on the exchange of carbon and water in an Amazonian rain forest”, *J. Geophys. Res.: Biogeosci.*, vol. 112, no. G3, pp. 1-16, 2007.

[10] Y. Yang, H. Guan, O. Batelaan, T.R. McVicar, D. Long, S. Piao, W. Liang, B. Liu, Z. Jin, and C.T. Simmons, “Contrasting responses of water use efficiency to drought across global terrestrial ecosystems”. *Sci. Rep.*, vol. 6, pp. 1-8, 2016.