

## NEAR-COLLISION SOLUTIONS ON NON-NEWTONIAN CENTRAL FORCE PROBLEM

**Saymon Santana**

National Institute for Space Research (LAC - INPE)  
Brazil  
saymonhss@gmail.com

**Elbert Macau**

National Institute for Space Research (LAC - INPE)  
Brazil  
elbert.macau@inpe.br

### Abstract

Central-force problems appear in various phenomena in science. This paper aims to study the near-collision solutions in Non-Newtonian central force.

Through a judicious change of variables we remove the singularity presented in the cases in which the solutions approach the origin. Through that, a fictitious flow about the singularity is created which allows the understanding of the near-collision dynamics.

### Key words

Central-force, collision, singularity.

### 1 Introduction

Several systems in physics are described by a vector field variant in time and or space. Let us take a particle moving in a force field  $\vec{F}$ .

Mathematically,  $\vec{F}$  is a vector belonging to  $\mathbb{R}^n$ . In a physical view,  $\vec{F}(\vec{X})$  can represent the force exerted on a particle located at  $\vec{X} \in \mathbb{R}^n$

The Newton's second law establishes the link between the physical and the mathematical concepts of force field. It asserts that, at any instant, a particle inserted in such force field moves in a such way that the force vector  $\vec{F}$  in a location  $\vec{X}$  of the particle is equal to the acceleration vector of the particle times his mass  $m$ , i.e.  $\vec{F} = m\vec{X}''$ .

This system of  $n$  second order differential equations may be written as a system of  $2n$  1st-order differential equations in  $\mathbb{R}^n \times \mathbb{R}^n$  such

$$\vec{X}'' = \vec{V} \quad (1)$$

$$\vec{V}' = \frac{1}{m}\vec{F}(\vec{X}) \quad (2)$$

where  $\vec{V}$  is the velocity of the particle.

The solution  $\vec{X}(t) \subset \mathbb{R}^n$  of 2nd-order system lies in a configuration space while the solution

$(\vec{X}(t), \vec{X}'(t)) \subset \mathbb{R}^n \times \mathbb{R}^n$  lies in the phase space of the system.

### 2 Central Force Systems

A force field  $\vec{F}$  is said to be a central type if  $\vec{F}(\vec{X})$  is a vector field points away or toward to every  $\vec{X}$ . Particularly, if the vector  $\vec{F}(\vec{X})$  is a scalar multiple of  $\vec{X}$ ,

$$\vec{F}(\vec{X}) = \lambda(\vec{X})\vec{X} \quad (3)$$

where the coefficient  $\lambda(\vec{X})$  depends on  $\vec{X}$ .

A central force model is often used to describes several phenomena in real world like a motion of an object orbiting around a massive body according to the *Newton's law of gravitation*, or the interaction between charged particles describer by the *Coulomb law*.

Since the force is conservative,  $\vec{F}$  can be written as a gradient of a potential scalar function  $U(\vec{X})$ , such that

$$\vec{F}(\vec{X}) = -\nabla U(\vec{X}) \quad (4)$$

In the special case known as a Newtonian central force system, the potential is of type

$$U(\vec{X}) = -\frac{1}{|\vec{X}|}$$

and the force varies with the inverse of the square distance.

### 3 Non-Newtonian central force problem

However, in some systems, such as those classical treated in [Buckingham, 1938], [Lennard-Jones, 1931], [Nicholson, 1989] and in many others related to Schrödinger's equation described in [Ikhdair and Sever, 2007], the scalar function varies in different ways from Newtonian cases.

Our focus is to discuss the classical cases of a Non-Newtonian central force problem. In those cases, the potential scalar function, which may represent an potential gravitational energy, is given by

$$U(\vec{X}) = \frac{1}{|\vec{X}|^\nu} \quad (5)$$

where  $\nu > 1$ .

In this case, the force

$$\vec{F} = \frac{\nu}{|\vec{X}|^{(\nu+1)}} \hat{X} \quad (6)$$

is not defined in  $\vec{X} = \vec{0}$ ; indeed the force becomes infinite as the moving particle approaches collision with the massive body.

Although the solutions in  $\vec{X} = \vec{0}$  may have not much importance in the real systems, it is very interesting to elucidate how solutions behave near to singularity. Our objective here is to remove this singularity by a judicious change of variables and time scalings.

Let us restrict our analysis to particles moving in plane ( $\vec{X} = (x, y) \in \mathbb{R}^2$ ), such that position can be written in polar coordinates (Fig. 1) as

$$(x, y) = (r \cos \theta, r \sin \theta)$$

and the velocity has components

$$(x', y') = (r' \cos \theta - r\theta' \sin \theta, r' \sin \theta + r\theta' \cos \theta).$$




Figure 1. Reference system for Central force problem

The solutions belong to configuration space  $\mathbb{C} = \mathbb{R}^2 - 0$ . Therefore, the phase space is denoted by  $\mathbb{P} = (\mathbb{R}^2 - 0) \times \mathbb{R}^2$ .

This phase space can be seen as a collection of all tangent vectors at each point  $\vec{X} \in \mathbb{C}$ . For a given  $\vec{X} \in \mathbb{R}^2 - 0$ , let  $\mathbb{T}_X = \{(\vec{X}, \vec{Y}) | \vec{Y} \in \mathbb{R}^2\}$ .  $\mathbb{T}_X$  is the tangent plane on the configuration space at  $\vec{X}$ .

Hence,  $\mathbb{P} = \bigcup_{\vec{X} \in \mathbb{C}} \mathbb{T}_X$  is the tangent space of the phase space.

We may introduce new variables  $(v_r, v_\theta)$  in the tangent space as

$$\vec{V} = v_r \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} + v_\theta \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix} \quad (7)$$

where  $v_r = r'$  and  $v_\theta = r\theta'$ .

Differentiating once more, we have

$$\begin{aligned} \vec{V}' &= \frac{\nu}{|\vec{X}|^{(\nu+1)}} \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} \\ &= \left(v_r' - \frac{v_\theta^2}{r}\right) \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} + \left(\frac{v_r v_\theta}{r} + v_\theta'\right) \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix} \end{aligned} \quad (8)$$

Rewriting the system in terms of the new coordinates

$$\begin{cases} r' = v_r \\ \theta' = \frac{v_\theta}{r} \\ v_r' = \frac{\nu}{r^{(\nu+1)}} + \frac{v_\theta^2}{r} \\ v_\theta' = -\frac{v_r v_\theta}{r} \end{cases} \quad (9)$$

Both the mechanical energy and angular momentum are integrals of motion, given respectively by

$$E = \frac{1}{2}(v_r^2 + v_\theta^2) + \frac{1}{r^\nu} \quad (10)$$

$$\vec{h} = r^2 \theta' (\hat{x} \times \hat{y}) \quad (11)$$

#### 4 Removing the singularity

In a similar way to what was done in [McGehee, 1974] we “blow up” the singularity via the following change of variables.

$$u_r = r^{\nu/2} v_r \quad (12)$$

$$u_\theta = r^{\nu/2} v_\theta \quad (13)$$

so that the system of equations becomes

$$\begin{cases} r' = r^{\nu/2} u_r \\ \theta' = r^{-(\frac{\nu}{2}+1)} u_\theta \\ u_r' = r^{-(\frac{\nu}{2}+1)} \left[ \frac{\nu u_r^2}{2} + u_\theta^2 + \nu \right] \\ u_\theta' = r^{-(\frac{\nu}{2}+1)} u_r u_\theta \left[ \frac{\nu}{2} - 1 \right] \end{cases} \quad (14)$$

If we multiply this equations by  $r^{(\frac{\nu}{2}+1)}$  we will remove the singularity that exists when  $r \rightarrow 0$ . In that case, the solution curve of the systems remain the same but are parameterized differently.

Strictly, we introduce a new time variable  $\tau$  such such that

$$\frac{dt}{d\tau} = r^{(\frac{\nu}{2}+1)} \quad (15)$$

In this new timescale the system becomes

$$\dot{r} = r^{\nu+1} u_r \quad (16)$$

$$\dot{\theta} = u_\theta \quad (17)$$

$$\dot{u}_r = \left[ \frac{\nu u_r^2}{2} + u_\theta^2 + \nu \right] \quad (18)$$

$$\dot{u}_\theta = u_r u_\theta \left[ \frac{\nu}{2} - 1 \right] \quad (19)$$

where the dot indicates differentiation with respect to  $\tau$ .

As can be seen, when  $r$  is small, the variation of  $\tau$  with respect to  $t$  is close to zero. Therefore, the scaled time  $\tau$  moves slowly then  $t$  near to the origin. In summary, the change of variables removes the singularity and allows to maintain the continuity of the solution.

## 5 Collision Solutions Analysis

In a precise manner, the singularity was replaced by a new set (well defined) given by  $r = 0$  and  $\theta$ , with  $u_r$  and  $u_\theta$  being arbitrary. Naturally, the set  $r = 0$  is an invariant set for the flow, provided that  $\dot{r} = 0$  when  $r = 0$ .

Since, in terms of new variables, the total energy relation is

$$\frac{1}{2}(v_r^2 + v_\theta^2) + 1 = hr^\nu \quad (20)$$

in  $r = 0$  we have a subset  $\Lambda$ , called *collision surface* and defined by

$$u_r^2 + u_\theta^2 = -2 \quad (21)$$

with  $\theta$  arbitrary, in which the solution behavior informs how the movement occurs in singularity's neighborhood.

This collision surface can be seen as a two-dimensional torus (Fig. 2), formed by a circle in  $\theta$  direction and another circle in the  $u_r u_\theta$  plane.




Figure 2. Collision surface  $\Lambda$

In this subspace, the system reduces to

$$\dot{\theta} = u_\theta \quad (22)$$

$$\dot{u}_r = u_\theta^2 \left[ 1 - \frac{\nu}{2} \right] \quad (23)$$

$$\dot{u}_\theta = u_r u_\theta \left[ \frac{\nu}{2} - 1 \right] \quad (24)$$

This system can be analyzed as follow:

- For  $\nu < 2$  (provided  $u_\theta \neq 0$ ) the coordinate  $u_r$  increases along any solution in  $\Lambda$ , as shown in Figure 3 for  $\nu = 1.99$  case.
- For  $\nu = 2$ , there is a bifurcation, in which  $u_r$  has no slope.
- For  $\nu > 2$ , the coordinate  $u_r$  decreases along any solution in  $\Lambda$  (since  $u_\theta \neq 0$ ), as shown in Figure 4 for  $\nu = 2.01$  case.




Figure 3. Phase plane for  $\nu = 1.99$




Figure 4. Phase plane for  $\nu = 2.01$

## 6 Final Remarks

This paper deals with an analytical way to remove the singularity existing in near collision solutions on Non-Newtonian potentials. Through a suitable change of variables, the system is parameterized and the previous divergent solution was replaced by a “smooth solution” when the system goes close to the singularity.

This type of artifice is extremely useful, especially in computational treatment of scattering problems, in which the analysis of dynamics close to singularity would produce an overflow, allowing analysis of configuration and phase spaces, even when the particle approaches the origin.

## Acknowledgements

The authors would like to thanks to CNPq (Process 435404/2016-4) and to Fapesp (grant 2015/50122-0 Sao Paulo Research Foundation) for the financial support.

## References

- Buckingham, R. A. (1938). The Classical Equation of State of Gaseous Helium, Neon and Argon *Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences* . 168 pp. 264-283 Wiley-Intersciences. New York.
- Ikhdaïr, S. M. Sever, R. (2007). Exact solutions of the radial Schrödinger equation for some physical potentials *Central European Journal of Physics*. Vol 5. N. 4, 516–527.
- LennardJones, A. F. (1931). Cohesion *Proceedings of the Physical Society*. Vol. 43, 461-482.
- McGehee, R. (1974). Triple collision in the collinear three body problem *Inventiones Mathematicae*. Vol. 27, p.191
- Nicholson, A. F. (1962). Bound States and Scattering in an r-2 Potential *Australian Journal of Physics*. Vol. 15, p.174